

Different approaches for constraining global climate models of the anthropogenic indirect aerosol effect

U. Lohmann

Institute for Atmospheric and Climate Science, ETH Zurich, Switzerland

J. Quaas, S. Kinne, J. Feichter

Max-Planck-Institute for Meteorology, Bundesstrasse 53, D-20146 Hamburg, Germany

Submitted to BAMS, Sept 29, 2005, revised June 14, 2006.

Short title:

Abstract. Assessments of the influence of aerosol emissions from human activities on the radiation budget, in particular via the modification of cloud properties, have been a challenge. In light of the variability to both aerosol properties and environmental properties affected by aerosols, observational evidence alone cannot provide accurate and global answers, because detailed observations are locally limited and/or lack statistical significance. Thus, our current understanding is predominantly derived from simulations with global models. General discrepancies to envelope (backward) modeling, however, suggest that many aerosol processes in global (forward) modeling are not properly considered. We recommend to use analytical derived parameterizations wherever possible. If an analytical method does not exist or is too demanding computationally, laboratory results augmented by field data are the second best approach. For the constraint of so-derived parameterizations at the GCM scale, we recommend to evaluate individual parameterizations using statistical relationships of satellite-retrieved quantities relevant to the process. The set of parameterizations may also be evaluated and improved using the data assimilation technique. To improve the quality of data-references to modeling, there is a need to link available atmospheric data from all scales, to establish and support validation networks and experiments, and a commitment to fine-tune and improve satellite retrievals in an iterative process even beyond the anticipated period of the mission. Only then more reliable estimates of the indirect aerosol effect can be expected.

1. Introduction

Aerosol particles resulting from human activity such as sulfate and carbonaceous aerosols have substantially increased the global mean aerosol burden since pre-industrial times. Aerosol particles can affect the climate system via several mechanisms. The most prominent impacts are (1) the reflection of solar radiation back to space (a 'direct' effect), (2) the absorption of solar radiation by soot and mineral dust to warm the atmospheric aerosol layer which could hinder cloud formation and/or cause cloud droplets to evaporate (a 'semi-direct' effect) and (3) the capability to act as condensation nuclei for (water and ice) clouds ('indirect effects'). The last effect, which is expected to increase the solar reflection of (water) clouds, is often distinguished into a cloud albedo and a cloud lifetime effect. The cloud albedo effect captures the process that polluted clouds with more but smaller droplets appear brighter (Twomey 1959), whereas the lifetime effect considers that polluted clouds with more but smaller droplets reduce the likelihood for cloud droplets to grow to raindrop size, thereby extending the cloud lifetime (Albrecht 1989). Modelling results suggest that these indirect effects are more important than the direct and semi-direct. Still despite many efforts large uncertainties remain for all simulated aerosol indirect effects (Penner et al. 2001; Ramaswamy et al. 2001).

Anderson et al. (2003) identified for the climatic impact of anthropogenic aerosols a discrepancy between climate model simulations and estimates from inverse models. Inverse models are conceptual models that derive the impact of aerosols by subtracting all other better quantified anthropogenic impacts from observed changes of surface temperatures where also the heat storage in oceans is considered (Knutti et al. 2002; Forest et al. 2002; Crutzen and Ramanathan 2003). Inverse methods constrain the aerosol impact on the energy balance at the top-of-the-atmosphere (TOA) to about -1 W m^{-2} , with an uncertainty range from 0 to -1.9 W m^{-2} (Anderson et al. 2003). In contrast, climate model simulations suggest a TOA forcing centered around -1.5 W

m^{-2} , and an uncertainty range that extends beyond -2.5 W m^{-2} (Anderson et al. 2003; Lohmann and Feichter 2005).

Thorough validation of aerosol-cloud interactions with observational data is missing in all climate model simulations of anthropogenic aerosol effects on clouds. There are some physical arguments in particular related to the treatment of the indirect effects, why estimates of the aerosol impact by current general circulation model (GCM) parameterizations suggest a stronger than expected cooling. There are a number of offsetting effects that are poorly, if at all considered in global modeling: (1) The semi-direct effect as mentioned above (Graßl 1979; Hansen et al. 1997); (2) the dispersion effect, which considers that the shape of the cloud droplet spectra is broader in polluted conditions (when the growth of the majority of the newly activated smaller aerosols is retarded (Liu and Daum 2002)), is believed to reduce both the albedo effect by 0.2 to 0.5 W m^{-2} (Peng and Lohmann 2003; Rotstayn and Liu 2003) and the cloud lifetime effect by a similar amount (Rotstayn and Liu 2005). (3) Anthropogenic aerosols can impact the microphysics of mixed-phase and ice clouds by acting as ice nuclei and thus accelerating ice-crystal induced precipitation, which would reduce cloud cover and lifetime (Lohmann 2002). (4) The more and smaller cloud droplets can also reduce the collision rate of snow crystals with cloud droplets (the riming process) in mixed-phase clouds (Borys et al. 2000 2003; Lohmann 2004). (5) More and smaller cloud droplets resulting from increasing aerosol concentrations suppress low-level rainout and aerosol washout. This allows transport of water and smoke to upper levels, thus elevating the onset of precipitation and the release the latent heat, which would result in more intensive convection (Rosenfeld and Woodley 2000; Andreae et al. 2004). These aerosol impacts on convective clouds have not been considered in climate modeling so far.

Most of these aerosol indirect effects have been deduced from the analysis of in-situ observations and satellite data on local scales. However, to be useful for global modeling with its coarse horizontal and vertical resolution, parameterizations must be developed,

that capture the essence of all aerosol-cloud interactions. And it must be demonstrated with available observational data that the developed parameterizations apply globally.

In chapter 2 we outline four different pathways to parameterizations. In chapter 3, methods to constrain parameterizations in global models are presented, and in chapter 4, we outline how those methods can be further developed in the future.

2. Methods to derive aerosol-cloud parameterizations

Processes, which act on spatial scales smaller than the horizontal (ca. 200×200 km²) and vertical resolution (ca. 20 tropospheric layers) and shorter than the typical time-step in global modeling (ca. 20 minutes), must be parameterized. It basically means to express variability on smaller scales in terms of large-scale variables. The concepts of the development of parameterizations can be divided into four different methods outlined below.

2.1 Derivation from first principles

If a process has an analytical solution, then this method should be used. Even if analytical solutions cannot be obtained directly, they often can be approximated with some simplifications. An example is the cloud droplet activation process, where particle size as a function of relative humidity is described by the Köhler curve for equilibrium conditions. Ghan et al. (1993) used the Köhler curve as the starting point for a parameterization of cloud droplet nucleation and assumed that the condensation rate can be related to the dry aerosol radius. This way droplet formation can be parameterized as a function of total aerosol number, vertical velocity and an activation parameter. Likewise, homogeneous ice crystal nucleation can be derived from theory if differences between the supersaturation at which freezing commences and the maximum supersaturation are neglected (Kärcher and Lohmann 2002).

Along the same lines the conversion of cloud droplets to form rain drops, the

autoconversion process, can be deduced analytically as discussed by Liu et al. (2004); Liu and Daum (2004); Liu et al. (2006). The autoconversion parameterizations traditionally requires a threshold cloud droplet size, above which the conversion to rain drops takes place. Instead of an arbitrary threshold, Liu et al. (2004) analytically derived an expression for this.

2.2 Derivation from laboratory studies

Laboratory data have been utilized in particular to study ice crystal formation. For instance, the effectiveness of mineral dust particles and/or black carbon particles to initiate contact or immersion freezing as a function of temperature is derived from a compilation of laboratory data (Diehl and Wurzler 2004; Diehl et al. 2006). These relationships have been applied in GCMs to study the importance of anthropogenic soot aerosols to serve as ice nuclei (Lohmann and Diehl 2006). This approach is promising because the relationship amongst variables of interest is studied in isolation. The derived parameterizations should be valid in the atmosphere as long as the heterogeneity and larger scales in real clouds do not influence these relationships. Thus, this method is the preferred method if no analytical solution is available.

2.3 Derivation from focused measurement campaigns

Field data at various continental and marine sites during cleaner and more polluted events have been summarized to derive robust relationships between (sub-micrometer size) aerosol number concentration (sometimes via the sulfate mass) and cloud droplet number concentration (Boucher and Lohmann 1995; Gultepe and Isaac 1996; Lin and Leaitch 1997). This concept was later extended for organic carbon and sea salt (Menon et al. 2002). These compiled data sets should be able to represent the spatial and temporal variability of droplet number concentration within a grid box of the model. Thus, they include all influences on the process in question, including those, that we

do not know. This is an advantage and a risk at the same time. In this regard, this method complements the laboratory method for processes that are more complex than can be studied in a laboratory setting. However, the sample size in a field experiment is normally not large enough to stratify these empirical data according to all influences in question. For example, empirical aerosol mass - cloud droplet number concentration relationships are limited with respect to the number and mixture of aerosol species that they take into account. Moreover, compilations of different data sets are complicated once different instruments are used for observing the same quantity.

2.4 Derivation from models with finer resolution

The cloud lifetime effect depends critically on the parameterization of the precipitation formation mechanism in warm clouds (Lohmann and Feichter 1997; Menon et al. 2002). These processes occur on much smaller scales. Thus, the growth of cloud droplets to precipitation size particles can be described by a stochastic (collection equation) relationship, as long as models keep track of different cloud droplet sizes. Following that idea, simulations of droplet growth with a cloud microphysical model have led to global model-suited parameterizations for the collision-coalescence process, the autoconversion rate and accretion rate (Khairoutdinov and Kogan 2000). This method is clearly not ideal, because it has no observational database. It is the last resort for a parameterization in the absence of any analytical solution or observations of the process in question. It is only useful if the GCM with different autoconversion schemes is validated against satellite data (Suzuki et al. 2004) so that the best scheme can be identified.

3. Constraints of parameterizations with large-scale observations

All methods to derive parameterizations outlined in the previous chapter have in common that they are based on theory or measurements valid for small scales and/or for specific situations. In a GCM the parameterizations need to be applied for any part of the globe, for any climate state and for the model's coarse spatial and temporal resolution. A widely used methodology to adapt parameterizations to changing environments and scales is to adjust (or 'tune') parameters to get a more 'realistic' match to available local data, e.g. (Rotstayn 2000). A better method would be to use observational data at adequate scales to infer these parameters. The use of information from satellite retrievals is particularly appealing, because they provide detailed cloud, aerosol, and radiation measurements at the large horizontal and temporal scales needed to evaluate GCMs. The resolution of the satellite goes down to scales able to distinguish between individual cloud-fields and cloud-free regions, which are of subgrid-scale for GCMs. There are, however, deficiencies in the use of satellite data in terms of temporal and vertical resolution, at least up to now.

3.1 Constraint with statistical relationships

Satellite-derived relationships can provide clues about the way specific parameterizations should work on the scales relevant in large-scale modeling. The advantage here is that statistical correlations from satellites are temporally and spatially more robust than individual measurements. This allows derivation of a desired quantity (e.g. cloud droplet number concentration) from predicted quantities (e.g. aerosol mass concentration). Correlations can also identify the overall effect in case that different effects with different signs and magnitudes are involved. Moreover, correlations analyze relative changes and therefore limitations to the absolute accuracy are acceptable.

Relationships are supposed to be valid also in a changing climate, whereas absolute values and currently measured distributions are not. For example, based on POLDER¹ satellite data it was found that the negative relationship between cloud droplet size and aerosol concentration is overestimated in global modeling (Lohmann and Lesins 2002). Based on similar findings, satellite derived relationships were used to revise existing parameterizations of cloud droplet activation from in-situ measurements used in GCMs (Quaas and Boucher 2005; Quaas et al. 2006).

3.2 Constraint using the data assimilation technique

The assimilation of observational data in global modeling is one approach to tie modeling to observations. Here, model parameters are relaxed towards observed values. Data assimilation can provide information of model deficiencies and measurement uncertainties especially in data rich areas. Assimilated climate models that require large model adjustments in certain areas or under certain conditions, offer obvious clues to poor representations of the adjusted model quantity. In turn model output of data assimilations can extend temporally or spatially sparse measurements. There is a European initiative GEMS that will follow this avenue as well as the US initiative PARAGON (Kahn et al. 2004). Moreover, this approach allows to test parameterizations, that were developed from the methods introduced in the previous chapter. Since the meteorology is prescribed for a particular time, testing could be as detailed as in numerical weather prediction. This enables model-evaluations around a specific events or at locations of dedicated measurement campaigns. Even then, we still face data issues because measurements (1) are usually unable to match the detail in modeling, (2) suffer from accuracy limitations and (3) may not be applicable at the temporal and spatial scales of modeling.

¹the acronyms are explained in the appendix

4. Recommendations

Table 1.

Table 2.

In order to constrain global models of the aerosol indirect effect, accurate and long-time observations for all relevant (aerosol, cloud, and environmental) quantities are required at adequate spatial and temporal resolutions (Tables 1 and 2). We should be able at least to address seasonal and interannual variability. Thus at a minimum data are needed for one complete annual cycle and if possible on a global or quasi-global (e.g. through intercalibrated ground networks such as AERONET) scale to establish links in space. Even longer records are certainly desirable given the year-to-year variability of aerosol (e.g. biomass burning) and of dynamics (e.g. El Nino, North Atlantic Oscillation) especially so, if data come from the same instrument (assuming that no instrument response changes and orbital changes occur).

Observations should discriminate the cloud type (shallow and deep convective, stratiform), cloud phase (water, ice, mixed-phase), the season, the region (arctic, mid-latitude, subtropic, and tropic), and the underlying surface (ocean, coastal, and continental). In terms of observational evidence on aerosol-cloud interactions we currently benefit from the increased capabilities of passive remote sensing (e.g. MODIS, MISR, PARASOL) and the establishment of passive and active remote sensing networks at surface sites (e.g. AERONET, BSRN, EARLINET). All of these measurements need to be combined in order to retrieve the whole suite of aerosol and cloud parameters that are relevant for the indirect aerosol effect. Instead of just comparing individual models with observations, Kinne et al. (2003 2006) pointed out the strength of investigating an ensemble of models at the same time not only to determine a most likely modeling value (model median) or regional model diversity but also to identify common biases in large-scale modeling.

More reliable correlations between aerosol, cloud and environmental atmospheric properties will be possible as available multi-year data-sets from the newer generation of satellite sensors are analyzed. Multi-spectral (MODIS, MERIS), multi-angle (MISR,

AATSR) and polarizing (PARASOL) sensors paired with soon to come vertical profiling (CALIPSO, CLOUDSAT) will allow for more accurate retrievals and especially provide detail for the interpretation of many observed correlations, such as to why MODIS data indicate a strong anti-correlation between aerosol optical depth and cloud top temperature (Koren et al. 2005). Even with the arrival of profiling sensors, one of the major shortcomings of aerosol-cloud correlations from space is the temporal resolution of polar-orbiting satellites, which is at best once per daytime.

In recognition that aerosol-cloud interactions (and in particular aerosol chemistry) occur on short time scales and even may display a diurnal cycle, there have been attempts to place capable sensors into the zero gravity point (L1) between sun and earth (e.g. the Triana project) or to improve capabilities on geostationary platforms (e.g. MSG). Their advantages and disadvantages are summarized in Table 3. Satellites in zero-gravity point and geostationary satellites deviate from nadir view at high latitudes and in case of the zero-gravity point satellites, deviations from nadir occur also towards sunrise and sunset. Global coverage and high-temporal resolution is preferred to address aerosol interactions. This could be reached either by a swarm of low cost time-delayed low-Earth orbiters or by at least 5 geostationary satellites preferably in combination with low-Earth orbit satellites. Initial insights on high temporally developing associations between aerosol and clouds although not globally are expected from the SEVIRI sensor (on MSG) data with 15 minutes-repeat capabilities on MSG (Meteosat) for studies mainly over Africa and the Atlantic. Particular interesting is a combined analysis with AERONET ground data, which provide more accuracy and detail on aerosol properties, at a similar temporal resolution.

Table 3.

However, even with these more sophisticated satellite instruments, retrieved radiances need to be converted into the relevant aerosol and cloud properties. The progress here is an iterative process that requires constant improvements of the retrievals based on careful comparisons to quality assured data, primarily from ground networks.

This emphasizes that any satellite mission needs to be complemented by a validation network of detailed field studies and long-term monitoring from the ground for the length of the satellite mission. A commitment even beyond the length of the mission is needed for staff to continuously compare, validate and reprocess the data. This certainly will be helped by an open data-policy.

In summary, we recommend to use analytical derived parameterizations wherever possible. If an analytical method does not exist or is too demanding computationally, laboratory results augmented by field data are the second best approach. For the constraint of so-derived parameterizations at the GCM scale, we recommend two complementary approaches. Individual parameterizations can be evaluated using statistical relationships of satellite-retrieved quantities relevant to the process. The set of parameterizations may also be evaluated and improved using the data assimilation technique making use of all available observational data.

Acknowledgments. This publication resulted from discussions held during the January, 2005 Specialty Conference on the Indirect Effect of Aerosols on Climate in Manchester, England, which was jointly sponsored by the International Global Atmospheric Chemistry Project (IGAC; a Core Project of the International Geosphere Biosphere Programme, IGBP) and by the U.S. NOAA and NASA programs. Financial support was also generously provided by the European Network of Excellence Atmospheric Composition Change, ACCENT.

Appendix: List of Acronyms:

AATSR: Advanced Along Track Scanning Radiometer

AERONET: AERosol RObotic Network

BSRN: Baseline Surface Radiation Network

CALIPSO: Cloud Aerosol Lidar and Infrared Pathfinder Satellite Observations

EARLINET: European Aerosol Research Lidar Network

GEMS: Global Earth system Monitoring using Space and in-situ data

GOES: Geostationary Operations Environmental Satellite

MERIS: MEidium Resolution Imaging Spectrometer

MISR: Multi-angle Imaging Spectro-Radiometer

MODIS: MODerate Resolution Imaging Spectro-radiometer

MSG: Meteosat Second Generation

PARAGON: Progressive Aerosol Retrieval and Assimilation Global Observing Network

PARASOL: Polarization and Anisotropy of Reflectances for Atmospheric Sciences coupled
with Observations from a Lidar

POLDER: POLarization and Directionality of the Earth's Reflectances

SEVIRI: Spinning Enhanced Visible and InfraRed Imager

References

- Albrecht, B. 1989. Aerosols, cloud microphysics, and fractional cloudiness. *Science* **245**, 1227–1230.
- Anderson, T. L., Charlson, R. J., Schwartz, S. E., Knutti, R., Boucher, O., Rodhe, H. and Heintzenberg, J. 2003. Climate forcing by aerosols - a hazy picture. *Science* **300**, 1103–1104.
- Andreae, M. O., Rosenfeld, D., Artaxo, P., Costa, A. A., Frank, G. P., Longo, K. M. and Silvas-Dias, M. A. F. 2004. Smoking rain clouds over the Amazon. *Science* **303**, 1337–1342.
- Borys, R. D., Lowenthal, D. H., Cohn, S. A. and Brown, W. O. J. 2003. Mountaintop and radar measurements of anthropogenic aerosol effects on snow growth and snowfall rate. *Geophys. Res. Lett.* **30**, doi: 10.1029/2002GL016,855.
- Borys, R. D., Lowenthal, D. H. and Mitchell, D. L. 2000. The relationships among cloud microphysics, chemistry, and precipitation rate in cold mountain cloud. *Atmos. Env.* **34**, 2593–2602.
- Boucher, O. and Lohmann, U. 1995. The sulfate-CCN-cloud albedo effect: A sensitivity study with two general circulation models. *Tellus Ser. B*, **47**, 281–300.
- Crutzen, P. J. and Ramanathan, V. 2003. The parasol effect on climate. *Science* **203**, 1679–1680.
- Diehl, K., Simmel, M. and Wurzler, S. 2006. Numerical simulations on the impact of aerosol properties and freezing modes on the glaciation, microphysics, and dynamics of convective clouds. *J. Geophys. Res.* **111**, doi:10.1029/2005JD005,884.
- Diehl, K. and Wurzler, S. 2004. Heterogeneous drop freezing in the immersion mode: Model calculations considering soluble and insoluble particles in the drops. *J. Atmos. Sci.* **61**, 2063–2072.
- Forest, C. E., Stone, P. H., Sokolev, A. P., Allen, M. R. and Webster, M. D. 2002. Quantifying

- uncertainties in climate system properties with the use of recent climate observations. *Science* **295**, 113–117.
- Ghan, S. J., Chuang, C. C. and Penner, J. E. 1993. A parameterization of cloud droplet nucleation, part i, single aerosol type. *Atmos. Res.* **30**, 197–221.
- Graßl, H. 1979. Possible changes of planetary albedo due to aerosol particles. In: W. Bach, J. Pankrath and W. Kellogg, eds., *Man's Impact on Climate*, Elsevier, New York.
- Gultepe, I. and Isaac, G. A. 1996. The relationship between cloud droplet and aerosol number concentrations for climate models. *Int. J. Climatol.* **16**, 1–6.
- Hansen, J., Sato, M. and Ruedy, R. 1997. Radiative forcing and climate response. *J. Geophys. Res.* **102**, 6831–6864.
- Kahn, R. A., Ogren, J. A., Ackerman, T. P., Bosenberg, J., Charlson, R. J., Diner, D. J., Holben, B. N., Menzies, R. T., Miller, M. A. and Seinfeld, J. H. 2004. Aerosol data sources and their roles within PARAGON. *Bulletin Of The American Meteorological Society* **85**, 1511–1522.
- Kärcher, B. and Lohmann, U. 2002. A parameterization of cirrus cloud formation: Homogeneous freezing of supercooled aerosols. *J. Geophys. Res.* **107**, doi: 10.1029/2001JD000,470.
- Khairoutdinov, M. and Kogan, Y. 2000. A new cloud physics parameterization in a large-eddy simulation model of marine stratocumulus. *Mon. Wea. Rev.* **128**, 229–243.
- Kinne, S., Lohmann, U., Feichter, J., Schulz, M., Timmreck, C., Ghan, S., Easter, R., Chin, M., Ginoux, P., Takemura, T., Tegen, I., Koch, D., Herzog, M., Penner, J., Pitari, G., Holben, B., Eck, T., Smirnov, A., Dubovik, O., Slutsker, I., Tanre, D., Torres, O., Mishchenko, M., Geogdzhayev, I., Chu, D. A. and Kaufman, Y. 2003. Monthly averages of aerosol properties: A global comparison among models, satellite data, and AERONET ground data. *J. Geophys. Res.* **108**, doi: 10.1029/2001JD001,253.
- Kinne, S., Schulz, M., Textor, C., Guibert, S., Balkanski, Y., Bauer, S., Berntsen, T., Berglen, T., Boucher, O., Chin, M., Collins, W., Dentener, F., Diehl, T., Easter, R., Feichter, J., Fillmore, D., Ghan, S., Ginoux, P., Gong, S., Grini, A., Hendricks, J.,

- Herzog, M., Horowitz, L., Isaksen, I., Iversen, T., Kirkevåg, A., Kloster, S., Koch, D., Kristjansson, J.-E., Krol, M., Lauer, A., Lamarque, J.-F., Lesins, G., Liu, X., Lohmann, U., Montanaro, V., Myhre, G., Penner, J., Pitari, G., Reddy, S., Seland, O., Stier, P., Takemura, T. and Tie, X. 2006. An AeroCom initial assessment - optical properties in aerosol component modules of global models. *Atmos. Chem. Phys.* **6**, 1815–1834.
- Knutti, R., Stocker, T. F., Joos, F. and Plattner, G.-K. 2002. Constraints on radiative forcing and future climate change from observations and climate model ensembles. *Nature* **416**, 719–723.
- Koren, I., Kaufman, Y. J., Rosenfeld, D., Remer, L. A. and Rudich, Y. 2005. Aerosol invigoration and restructuring of atlantic convective cloudus. *Geophys. Res. Lett.* **32**, doi:10.1029/2005GL023,187.
- Lin, H. and Leitch, W. R. 1997. Development of an in-cloud aerosol activation parameterization for climate modelling. In: *Proceedings of the WMO Workshop on Measurement of Cloud Properties for Forecasts of Weather, Air Quality and Climate*. World Meteorol. Organ., Geneva, 328–335.
- Liu, Y. and Daum, P. H. 2002. Indirect warming effect from dispersion forcing. *Nature* **419**, 580–581.
- Liu, Y., Daum, P. H. and McGraw, R. 2004. An analytical expression for predicting the critical radius in the autoconversion parameterization. *Geophys. Res. Lett.* **31**, doi: 10.1029/2003GL019,117.
- Liu, Y. G. and Daum, P. H. 2004. Parameterization of the autoconversion process. part i: Analytical formulation of the kessler-type parameterizations. *Journal of the Atmospheric Sciences* **61**, 1539–1548.
- Liu, Y. G., Daum, P. H., McGraw, R. and Wood, R. 2006. Parameterization of the autoconversion process. part ii: Generalization of sundqvist-type parameterizations. *Journal of the Atmospheric Sciences* **63**, 1103–1109.

- Lohmann, U. 2002. A glaciation indirect aerosol effect caused by soot aerosols. *Geophys. Res. Lett.* **29**, doi: 10.1029/2001GL014,357.
- Lohmann, U. 2004. Can anthropogenic aerosols decrease the snowfall rate? *J. Atmos. Sci.* **61**, 2457–2468.
- Lohmann, U. and Diehl, K. 2006. Sensitivity studies of the importance of dust ice nuclei for the indirect aerosol effect on stratiform mixed-phase clouds. *J. Atmos. Sci.* **63**, 968–982.
- Lohmann, U. and Feichter, J. 1997. Impact of sulfate aerosols on albedo and lifetime of clouds: A sensitivity study with the ECHAM GCM. *J. Geophys. Res.* **102**, 13,685–13,700.
- Lohmann, U. and Feichter, J. 2005. Global indirect aerosol effects: A review. *Atmos. Chem. Phys.* **5**, 715–737.
- Lohmann, U. and Lesins, G. 2002. Stronger constraints on the anthropogenic indirect aerosol effect. *Science* **298**, 1012–1016.
- Menon, S., DelGenio, A. D., Koch, D. and Tselioudis, G. 2002. GCM simulations of the aerosol indirect effect: Sensitivity to cloud parameterization and aerosol burden. *J. Atmos. Sci.* **59**, 692–713.
- Peng, Y. and Lohmann, U. 2003. Sensitivity study of the spectral dispersion of the cloud droplet size distribution on the indirect aerosol effect. *Geophys. Res. Lett.* **30**, doi: 10.1029/2003GL017,192.
- Penner, J. E., Andreae, M., Annegarn, H., Barrie, L., Feichter, J., Hegg, D., Jayaraman, Leaitch, R., Murphy, D., Nganga, J. and Pitari, G. 2001. Aerosols, their direct and indirect effects. In: J. T. Houghton, Y. Ding, D. J. Griggs, M. Noguer, P. J. van der Linden, X. Dai, K. Maskell and C. A. Johnson, (Eds.), eds., *Climate Change 2001: The Scientific Basis. Contribution of working group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge Univ. Press, New York. 881.
- Quaas, J. and Boucher, O. 2005. Constraining the first aerosol indirect radiative forcing in the LMDZ GCM using POLDER and MODIS satellite data. *Geophys. Res. Lett.* **32**, doi: 10.1029/2005GL023,850.

- Quaas, J., Boucher, O. and Lohmann, U. 2006. A new estimate of the aerosol indirect radiative forcing by constraints of global climate models using satellite datasets. *Atmos. Chem. Phys.* **6**, 947 – 955.
- Ramaswamy, V., Boucher, O., Haigh, J., Hauglustaine, D., Haywood, J., Myhre, G., Nakajima, T., Shi, G. Y. and Solomon, S. 2001. Radiative forcing of climate change. In: J. T. Houghton, Y. Ding, D. J. Griggs, M. Noguer, P. J. van der Linden, X. Dai, K. Maskell and C. A. Johnson, eds., *Climate Change 2001: The Scientific Basis. Contribution of working group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge Univ. Press, New York. 349–416.
- Rosenfeld, D. and Woodley, W. L. 2000. Deep convective clouds with sustained supercooled liquid water down to -37.5°C . *Nature* **405**, 440–442.
- Rotstayn, L. D. 2000. On the "tuning" of autoconversion parameterizations in climate models. *J. Geophys. Res.* **105**, 15,495–15,508.
- Rotstayn, L. D. and Liu, Y. 2003. Sensitivity of the first indirect aerosol effect to an increase of cloud droplet spectral dispersion with droplet number concentration. *J. Climate* **16**, 3476–3481.
- Rotstayn, L. D. and Liu, Y. 2005. A smaller global estimate of the second indirect aerosol effect. *Geophys. Res. Lett.* **32**, doi:10.1029/2004GL021,922.
- Suzuki, K., Nakajima, T., Numaguti, A., Takemura, T., Kawamoto, K. and Higurashi, A. 2004. A study of the aerosol effect on a cloud field with simultaneous use of GCM modeling and satellite observations. *J. Atmos. Sci.* **61**, 179–194.
- Twomey, S. A. 1959. The nuclei of natural cloud formation. Part II: The supersaturation in natural clouds and the variation of cloud droplet concentrations. *Geofis. Pure Appl.* **43**, 227–242.

Received _____

With the extension package 'AGU++' by P.W.Daly, version 1.5 from 1996/08/20

Tables

Table 1. Physical parameters related to aerosols, clouds and dynamics needed to evaluate parameterizations of aerosol indirect effects.

Parameter	description	typical value or range
N_a	Aerosol number concentration	$10^6 - 10^{11} m^{-3}$
SD_a	Aerosol size distribution	log-normal
SF_a	Soluble fraction of aerosol population	0 - 1
AT	Aerosol type	-
SSA	Single scattering albedo	0.6 - 0.99
BC(z)	Position of black carbon w.r.t. the cloud	above/in/below
CC	Cloud cover	0 - 1
N_d	Cloud droplet number concentration	$10^6 - 10^{10} m^{-3}$
N_i	Ice crystal number concentration	$10^2 - 10^8 m^{-3}$
SD_d	Cloud droplet size distribution	log-normal or gamma distribution
SD_i	Ice crystal size distribution	log-normal or gamma distribution
LWC	Liquid water content	$0 - 10^{-3} kg m^{-3}$
IWC	Ice water content	$0 - 10^{-3} kg m^{-3}$
AU	Autoconversion rate	$0 - 10^{-6} kg kg^{-1} s^{-1}$
w	Vertical wind speed	-10 to 10 $m s^{-1}$
RH	Relative humidity	20 - 100% (up to 170% w.r.t. ice)
TKE	Turbulent kinetic energy	$0 - 10 m^2 s^{-2}$
T	temperature	210 - 273 K

Table 2. Attribution of the parameters listed in Table 1 to the evaluation of the different aerosol indirect effect mechanisms, including the required data resolution.

Evaluation of	aerosols	clouds	large-scale environment	required resolution (x, z, t)
Cloud albedo effect	N_a , SD_a , SF_a , AT	N_d , SD_d , LWC, CC	w, RH	1km, 100m, 1h
Cloud lifetime effect	-	N_d , SD_d , LWC, AU, CC	TKE	1km, 100m, 1h
Semi-direct effect	N_a , SSA, BC(z)	LWC, CC	RH	10km, 1km, 6h
Aerosol effects on mixed-phase and ice clouds	N_a , SD_a , SF_a , AT	N_i , SD_i , IWC, CC	w, RH, T	1km, 100m, 1h

Table 3. Advantages and disadvantages of the different satellite observing concepts.

Satellite-orbits	LEO ¹	GEO ²	L1 ³ /L2 ⁴
Altitude above ground	low	mid	high
Global coverage	yes	no	yes
Temporal resolution	low	high	high
Overpass time	specific time	always	day(L1), night(L2)
High latitudes	good	poor	poor
Examples	MODIS	GOES	TRIANA (planned)

¹LEO: low-Earth orbit (e.g. polar orbiting - crossing up and down the equator)

²GEO: geostationary Earth orbit (placed at a fixed longitude over the equator)

³L1: placed in the equal attraction point of Earth and Sun (between Earth and Sun)

⁴L2: placed in the equal attraction point of Earth and Sun (behind the Earth). At the Lagrangian points L1 or L2 a satellite can 'move with the sun'